

Bibliography

Publications using the continual trials approach to spontaneous recognition tasks

Ameen-Ali, K.E., Eacott, M.J. & Easton, A. (2012). A new behavioural apparatus to reduce animal numbers in multiple types of spontaneous object recognition paradigms in rats. *Journal of neuroscience methods* **211**(1): 66-76.

Seel, S.V., Eacott, M.J., Langston, R.F. & **Easton, A.** (2017). Cholinergic input to the hippocampus is not required for a model of episodic memory in the rat, even with multiple consecutive events. *Behavioural Brain Research* (10.1016/j.bbr.2017.06.001)

Review paper summarising the advantages of moving beyond standard approaches to spontaneous recognition tasks

Ameen-Ali, K.E., Easton, A. & Eacott, M.J. (2015). Moving beyond standard procedures to assess spontaneous recognition memory. *Neuroscience and Biobehavioral Reviews* **53**: 37-51.

Uses of an episodic spontaneous recognition task (one trial a day)

Davis, K.E., **Easton, A., Eacott, M.J.** & Gigg, J. (2013). Episodic-Like Memory for What-Where-Which Occasion is Selectively Impaired in the 3xTgAD Mouse Model of Alzheimer's Disease. *Journal of Alzheimer's Disease* **33**(3): 681-698.

Davis, K.E., **Eacott, M.J., Easton, A.** & Gigg, J. (2013). Episodic-like memory is sensitive to both Alzheimer's-like pathological accumulation and normal ageing processes in mice. *Behavioural Brain Research* **254**: 73-82.

Easton, A., Fitchett, A., Eacott, M.J. & Baxter, M.G. (2011). Medial septal cholinergic neurons are necessary for context-place memory but not episodic-like memory. *Hippocampus* **21**(9): 1021-1027.